

VERÖFFENTLICHUNGEN 2012

Abt. Pflanzenzüchtung

- Cleemput, S. and H. Becker, 2012: Genetic variation in leaf and stem glucosinolates in resynthesized lines of winter rapeseed (*Brassica napus* L.). *Genet Resour Crop Evol* 59, 539-546.
- Dimov, Z., E. Suprianto, F. Hermann and C. Möllers, 2012: Genetic variation for seed hull and fibre content in a collection of European winter oilseed rape material (*Brassica napus* L.) and development of NIRS calibrations. *Plant Breeding* 131, 361-368.
- Duc, G., W. Link, P. Marget, F.L. Stoddard, A.M. Torres and J.I. Cubero, 2011: Genetic Adjustment to Changing Climates: Faba Bean. Chapter 10. Pp. 269-286. In: Yadav, S. S. et al. (Eds.): *Crop Adaptation to Climate Change*. Wiley-Blackwell, Chichester, West Sussex, UK. ISBN: 978-0-8138-2016-3.
- Flores, F., S. Nadal, I. Solis, J. Winkler, O. Sass, F.L. Stoddard, W. Link, B. Raffiot, F. Muel and D. Rubiales, 2012: Faba bean adaptation to autumn sowing under European climates. *Agron. Sustain. Dev.* 32, 727-734.
- Girke, A., A. Schierholt and H.C. Becker, 2012: Extending the rapeseed gene pool with resynthesized *Brassica napus* L. I. Genetic diversity. *Genet Resour Crop Evol*, 59, 1441-1447.
- Girke, A., A. Schierholt and H.C. Becker, 2012: Extending the rapeseed gene pool with resynthesized *Brassica napus*. II. Heterosis. *Theor Appl Genet* 124, 1017-1026.
- Gruber, S., A. Hüskens, A. Dietz-Pfeilstetter, C. Möllers, E. A. Weber, F. Stockmann, H. Thöle, J. Schatzki, K. Dowideit, M. Renard, H.C. Becker, J. Schiemann and W. Claupein, 2012: Biological confinement strategies for seed- and pollen-mediated gene flow of GM canola (*Brassica napus* L.). *AgBioForum* 15, 44-53.
- Horneburg, B. and J.R. Myers, 2012: Tomato: Breeding for improved disease resistance in fresh market and home garden varieties. In: Lammerts van Bueren, E.T. and J.R. Myers (Eds.). *Organic Crop Breeding*. Chapter 15. Pp. 239-250. Wiley-Blackwell, Chichester, West Sussex, UK. ISBN: 978-0-470-95858-2.
- Jauker, F., B. Bondarenko, H. C. Becker and I. Steffan-Dewenter, 2012: Pollination efficiency of wild bees and hoverflies provided to oilseed rape. *Agricultural and Forest Entomology*. 14, 81-87.
- Kessel, B., A. Schierholt and H. C. Becker, 2012: Nitrogen use efficiency in a genetically diverse set of winter oilseed rape (*Brassica napus* L.). *Crop Science* 52, 2546-2554.
- Klaedtke S. and B. Horneburg, 2012: Aromatische Tomatensorten trotz Krautfäule - auch in Luxemburg? *Gaart an Heem*. Ausgabe Mai 2012, S. 132.
- Link, W. and L. Ghaouti, 2012: Faba Bean: Breeding for organic farming systems. In: Lammerts van Bueren, E.T. and J.R. Myers (Eds.): *Organic Crop Breeding*. Chapter 13. Pp. 215-226. Wiley-Blackwell, Chichester, West Sussex, UK. ISBN: 978-0-470-95858-2.
- Ofori, A., A. Schierholt and H.C. Becker, 2012: Biomass yield and heterosis of crosses within and between European winter cultivars of turnip rape (*Brassica rapa* L.). *J Appl Genetics*, 53, 31-35.

Rudolphi, S., H.C. Becker, A. Schierholt and S. von Witzke-Ehbrecht, 2012: Improved estimation of oil, linoleic and oleic acid and seed hull fractions in safflower by NIRS. J Am Oil Chem Soc. 89, 363 -369.

Zamani-Nour, S., R. Clemens and C. Möllers, 2012: Cytoplasmic diversity of *Brassica napus* L., *Brassica oleracea* L. and *Brassica rapa* L. as determined by chloroplast microsatellite markers. Genet Resour Crop Evol. DOI: 10.1007/s10722-012-9891-x.

Monographien

Schatzki, J., 2012: Genetic variation and inheritance of secondary seed dormancy in winter oilseed rape (*Brassica napus* L.). Diss. Universität Göttingen. 84 S.
<http://webdoc.sub.gwdg.de/diss/2012/schatzki>